Caries Experience and Treatment Needs Among Residents of an Internally Displaced Persons (IDPs) Camp in Edo State, Nigeria.

*Okuonghae OE, **Akhigbemen O, ***Onagaumah JJ, **Udobi L, **Enabulele JE, ***Azodo CC

*Department of Preventive Dentistry, University of Benin Teaching Hospital **Department of Restorative Dentistry, University of Benin Teaching Hospital ***Department of Periodontics, University of Benin Teaching Hospital ABSTRACT

Correspondence
Dr. O. E Okuonghae
Department of Preventive Dentistry, University
of Benin Teaching Hospital
Benin City
Nigeria.
Email: elosticker@gmail.com

Citation: Okuonghae OE, Akhigbemen O, Onagaumah JJ, Udobi L, Enabulele JE, Azodo CC (2024). Caries Experience and Treatment Needs among Residents of an Internally Displaced Persons (IDPs) Camp in Edo State, Nigeria. Nig J Res Orofac Infect Oncol; 7(1&2):c1-c2.

INTRODUCTION

The act of terrorism has become persistent in the Northern part of Nigeria resulting in significant impact in various aspects of the economy and one of such is the displaced people crisis within the country resulting in persons referred to as Internally displaced persons (IDPs) (Nnawuihe & Otakhoigbogie, 2021; Arheiam et al., 2023). These persons are considered vulnerable as they are at a higher risk of disparity in oral health care and bear a disproportionate burden of oral diseases. One of the most prevalent diseases of the oral cavity is dental caries, which when untreated can cause pain, swelling and facial infections, among other consequences which can affect the overall quality of life (Razan et al., 2022). There is a need to assess the caries experience, treatment need and dental caries severity among these groups. The objective of the study was to evaluate the prevalence of dental caries, the treatment needs, and the dental caries severity among internally displaced persons living in a camp.

METHODS

An analysis of secondary data obtained at an oral health outreach carried out among adults and children residing in an Internally Displaced Persons' Camp at Uhogua village forest reserve, Edo State, Nigeria as part of the World Oral Health Day by the Edo state branch of Nigeria Dental Association. Data retrieved were age, gender, decayed teeth, missing, and treatment done. The prevalence, severity, and treatment need with regard to dental caries were calculated before and after intervention. IBM SPSS version 26.0 was used for descriptive and inferential analysis at 95% confidence interval with a p-value set at < 0.05 significance.

RESULTS

A total of 83 patient records were retrieved with an age range of 3-43 and a mean age of 15.01± 6.9. with the majority within the age group of 10-20years (65.1%). Male had a slightly higher representation of 43 (51.8%) with a male-female ratio of 1.1:1.

Caries prevalence was 33.7% with females having higher prevalence (53.9%). The mean Decayed, Missing, and

Filled Teeth (mDMFT) for adults was 0.45, while the mean dmft for children with primary teeth was 0.44. The mean dmft in females (0.55) was statistically significantly higher than that recorded in males (0.27). The treatment need was calculated before and after intervention with the mean treatment need before intervention 32.5± 5.2 and the mean treatment need after intervention 20.7± 4.4 showing a slight reduction in treatment need after intervention.

CONCLUSION

The study shows that the IDPs had a dental caries prevalence of 33.7% suggesting that persons living in IDP camps need more attention to dental health services. Recommended strategies to further reduce the dental caries prevalence and experience will be instituting oral health education programs and the creation of dental safety net clinics at the camp which will increase access to preventive oral health services for this group. The IDPs benefited from free oral health-care services.

Financial support and sponsorship

This work received no specific grant from any funding agency in the public, commercial or not-for-profit sectors

Conflicts of interest

The authors declare that they have no conflicts of interest.

REFERENCES

- Arheiam A, Alhashani A, Kwidir T, Bosif Y, Ballo L, Tantawi ME (2023]). Untreated dental caries among Libyan children during and after the war and in internally displaced person camps. Community Dent Oral Epidemiol;51(4):636-643.
- Nnawuihe UC, Otakhoigbogie U (2021). Oral disease burden amongst residents of an internally displaced persons camp in Nigeria. Nig J Dent Res.; 6(2):109-115.
- Razan OT, Nada HM, Isam MA, Elhadi MA (2022). Oral Health Status among Refugees and Internally Displaced Children (3-11 Years Old) from Western Kordofan State, Sudan. Saudi J Oral Dent Res.;7(5): 142-146
- Umeizudike KA, Dedeke AA, Nzomiwu CL, Ekowmenhenhen UI (2019). Oral health status and treatment needs of internally displaced persons. Sahel Med J; 22:207-13